

X-ray sources in globular clusters

Frank Verbunt

Astronomical Institute, Postbox 80.000, 3508 TA Utrecht, the Netherlands

Abstract. Observations with BeppoSAX, RXTE and Chandra suggest that many of the bright X-ray sources in globular clusters have ultrashort binary periods. This is remarkable as such systems are not easily formed. With accurate optical astrometry of HST images, the large numbers of low-luminosity X-ray sources discovered with Chandra can be classified as quiescent low-mass X-ray binaries, pulsars, cataclysmic variables, and magnetically active binaries. The number of cataclysmic variables is found to scale with the number of close stellar encounters.

INTRODUCTION

The known populations of X-ray sources in globular clusters have grown with the increasing sensitivity of X-ray detectors, from bright low-mass X-ray binaries discovered in the 1970s (UHURU, OSO-7, Ariel-V), to faint low-mass X-ray binaries and cataclysmic variables in the 1980s (Einstein, HEAO-2), recycled radio pulsars in the 1990s (ASCA, ROSAT), and magnetically active close binaries in the 2000s (Chandra). In this review I summarize the new knowledge gained in the last years about the bright X-ray sources, and discuss in particular the possibility that many of these have ultrashort orbital periods (Sect. 2). Then I explain how various new technical possibilities of Chandra, XMM and HST expand our knowledge of the faint sources (Sect. 3). Some considerations on formation rates end the paper.

THE BRIGHT SOURCES: NEUTRON STARS ACCRETING AT HIGH RATES

We now know 13 bright X-ray sources ($L_{0.5-2.5\text{keV}} > 10^{35.5} \text{ erg s}^{-1}$, say) in 12 globular clusters of our Galaxy (Table 1). Most were discovered in the 1970s with the first X-ray satellites; relatively recent additions are the source in Terzan 6, discovered in the ROSAT All Sky Survey, and the two sources into which the bright X-ray emission from NGC 7078 – known since the 1970s – was resolved by Chandra. X-ray bursts were detected from some sources soon after discovery, and some sources (Terzan 1 and Terzan 5) were initially detected only during bursts. It required the large temporal and spatial coverage by the BeppoSAX Wide Field Cameras and the RXTE All Sky Monitor to find that all but one of the bright sources in globular clusters are X-ray bursters, and thus are neutron stars accreting from a companion. The 13th source, in NGC7078, may be a neutron star as well.

TABLE 1. List of the thirteen bright X-ray sources in globular clusters with the year of discovery, year of the first observation of a burst, absolute magnitude in B (or U, J, as indicated), orbital period, and indications whether the source is a transient (T), and whether the source has properties associated with an ultrashort period (U) or with a normal period (N) in its optical magnitude (O), X-ray spectrum (X), or burst maximum (B). See text for explanation. Detailed references are given in Verbunt & Lewin (2004).
^aLuminous X-ray emission from NGC 7078 was already found in 1974. ^bor its alias at 13.2 m.

cluster	disc	burst	M_λ	P_b	TOXB
NGC 1851	1975	1976	5.6B		UUU
NGC 6440	1975	1999	3.7B		T–N–
NGC 6441	1974	1987	2.4B	5.7hr	–NN
NGC 6624	1974	1976	3.0B	11.4m	UUU
NGC 6652	1985	1998	5.6B		UUU
NGC 6712	1976	1980	4.5B	20.6m ^b	UUU
NGC 7078-1 ^a	2001		0.7B	17.1hr	---
NGC 7078-2 ^a	2001	1990	3.1U		--U
Terzan 1	1981	1981			T---
Terzan 2	1977	1977			–NU
Terzan 5	1981	1981	1.7J		T–U–
Terzan 6	1991	2003		12.36h	T–N–
Liller 1	1976	1978			T---

Remarkably, of the five orbital periods currently known, two are ultrashort, at periods of 11.4 m and 20.6 m (or its alias 13.2 m). I return to this below. Five of the 13 sources are transients, in the sense that their luminosity at times has dropped below $\sim 10^{34}$ erg s^{–1}.

Recently, two remarkable illustrations of the danger of identifications without accurate positions have been found. The first is the discovery that the bright ultraviolet source in NGC 6652, with a period of 43.6 m (Deutsch et al. 2000), is *not* the optical counterpart of the bright X-ray source in this cluster, but of a second, relatively faint X-ray source, presumably a faint low-mass X-ray binary (Heinke et al. 2001). The second is the discovery that the single faint source significantly detected in a Chandra observation of Terzan 1 is *not* the quiescent counterpart of the transient in this cluster, but a different source (see Figure 1). A warning that the optical centers of globular clusters are not always as accurate as advertised is provided by the study of Terzan 6: the hitherto assumed optical center was probably offset from the actual center by the influence of a bright star outside the cluster. A re-determination of the cluster center, in which each star gets the same weight irrespective of its brightness, places the center very close to the accurately determined position of the X-ray source (in ’t Zand et al. 2003).

With three ROSAT observations of NGC 6624, van der Klis et al. (1993) showed that the 11.4 m period of the X-ray source in this cluster decreases, in contradiction of the model in which the donor delivering matter to the neutron star is a white dwarf. The negative period derivative is confirmed by Chou & Grindlay (2001). Some hesitation in accepting this derivative as intrinsic may remain, since the differences between observed periods as a function of time and a quadratic fit look irregular. Also, it is possible that acceleration of the binary in the cluster potential affects its observed period derivative: the optical counterpart of the X-ray source is very close to the cluster center ($0.6 \pm 0.3''$, King et al. 1993). Nonetheless, one must now seriously consider the possibility that the 11.4 m period is decreasing rather than increasing.

Are there models which can explain this? To discuss this, we note first that the




FIGURE 1. *Left: Accurate positions for the bright source in Terzan 1 were determined with EXOSAT (× with dashed error circle, from Parmar et al. 1989) and with three independent ROSAT observations (+ with solid error circles, redetermined by the author after Johnston et al. 1995). Coordinates are with respect to the cluster center. The bright source had switched off in 1999 (Guainazzi et al. 1999). The position of the one source detectable in a short Chandra observation, indicated as the small error circle on the right, is not compatible with the position of the previously bright source (Wijnands et al. 2002). Right: Comparison of the X-ray spectrum of a quiescent low-mass X-ray binary (NGC6205 Ga, below left) and a radio pulsar (in M28, up right). Note the sensitivity of ROSAT to very soft photons. Data from ROSAT and XMM from Gendre et al. (2003); from Chandra from Becker et al. (2003).*

formation of the bright X-ray binaries in globular clusters is thought to follow two main paths: either a tidal capture of a closely passing star by a single neutron star, or an exchange encounter in which a neutron star encounters a binary and swaps place with one binary star. If a low-mass main-sequence star is captured, and starts transferring mass to the neutron star, the orbit will decrease until the donor star becomes degenerate, after which the orbit expands again. The minimum period reached this way is near 70-80 m, and shorter periods cannot be explained this way. If a more massive main-sequence star is captured, and evolves into a giant before mass transfer starts, the mass transfer is unstable and a spiral-in of the neutron star into the envelope of the giant may bring it into a close orbit with the core of the giant. This must have happened in the past, when stars with a mass sufficiently high to induce unstable mass transfer still existed in globular clusters: it may take several billion years before the core of the giant, cooled into a white dwarf, comes into contact under the influence of gravitational radiation and starts mass transfer. Such ultrashort period binaries can therefore be observed today (Rasio et al. 2000). If a neutron star collides directly with a giant, the envelope will be expelled, and the core may enter an orbit around the neutron star. A white-dwarf donor of mass M_c fills its Roche-lobe at an orbital period of roughly $P_b \sim 1^m (M_\odot/M_c)$; thus ultrashort periods are possible, but the period must lengthen with time as the mass of the white dwarf decreases (Verbunt 1987). Tutukov et al. (1985) note that ultrashort periods can also be reached in binaries in which a star starts to transfer mass to its companion after it has already evolved somewhat, and depleted its core from hydrogen. The donor




FIGURE 2. Left: Absolute B magnitude of optical counterparts to X-ray sources in globular clusters as a function of the X-ray luminosity (expressed in Eddington luminosity $2 \times 10^{38} \text{ erg s}^{-1}$) and orbital period (in hours). Data from Deutsch et al. (2000) and Heinke et al. (2001), see Table 1. The solid and dashed lines give the average relation and spread around the average for X-ray binaries in the galactic disk, as derived by Van Paradijs & McClintock (1994). Right: maximum observed luminosities L_m during X-ray bursts, compared with the Eddington limits for hydrogen-rich ($X = 0.73$) and hydrogen-poor ($X = 0$) matter (lower and upper gray band, respectively). After Kuulkers et al. (2003). In both figures: $\square$ normal period, $\triangle$ ultrashort period, $\circ$ period not known (set at 1 hr for the left plot). A filled symbol indicates that the peak flux was determined for an ordinary burst, and may be lower than the Eddington limit.

then becomes degenerate at smaller mass, and shorter orbital period. Tutukov et al. do a limited number of calculations and find a minimum period of about 20 m; Podsiadlowski et al. (2002) show that periods as short as 5 m can be reached this way. This scenario may be called 'magnetic capture'.

Van der Sluys et al. (these proceedings, see also Van der Sluys et al. 2004) note that none of the relevant computations of magnetic capture done by Pylyser & Savonije (1988) reach such extremely short periods, and investigate in some detail which initial conditions (initial binary period and donor mass) lead to ultrashort orbital periods *within a Hubble time*. They find that only binaries in a very small range of initial orbital periods and in a small range of donor masses evolve to ultrashort periods within the Hubble time, and that each of these binaries spends only a small fraction of its evolution at ultrashort periods. They conclude that only an extremely small fraction (< 0.001) of a population of X-ray binaries at any time can be expected to be at ultrashort periods. Whereas the observation of one such system amongst the thirteen globular cluster sources could be explained by magnetic capture, the observation of two or more demands another explanation.

In this context it is worthy of note that there are good indications that many of the bright X-ray sources in globular clusters have ultrashort periods, as first pointed out by Deutsch et al. (1996). As seen in Table 1, two of five orbital periods known are ultrashort. Indirect evidence for ultrashort periods is provided by three observations. First, it is known that the optical light of bright low-mass X-ray binaries is dominated




FIGURE 3. Fitted values of model parameters of X-ray spectra of X-ray sources in globular clusters. Shown are the blackbody temperatures T_{in} of the photons leaving the inner disk as a function of the projected inner disk radius $R_{in}\sqrt{\cos(i)}$ (left) and of the blackbody temperatures T_o of photons entering the Comptonization region (right). After Sidoli et al. (2001); NGC 6652 added from Parmar et al. 2001, Terzan 5 added from Heinke et al. (2003). Symbols as in Fig. 2.

by reprocessing of X-rays that impinge on the accretion disk. Thus, if the disk is small and/or if the X-ray luminosity is small, one expects a low optical luminosity. Van Paradijs & McClintock (1994) derive that the expected absolute visual magnitude scales with disk size (via orbital period P_b) and X-ray luminosity L_x roughly as $M_V \propto P_b^{2/3} L_x^{1/2}$. Figure 2 shows that the transition between systems with normal periods (i.e. periods above 80 m) and those with ultrashort periods lies near $M_B = 3$. The B magnitudes of the sources in NGC 1851 and NGC 6652 thus suggest ultrashort periods.

The second indication is the maximum luminosity reached in a burst. Kuulkers et al. (2003) have analyzed an extensive data set of X-ray bursts of sources in globular clusters, and note that the maximum luminosity reached in radius expansion bursts, when the source is assumed to reach the Eddington limit, corresponds to the Eddington limit for hydrogen-poor material in the two systems with ultrashort periods, and to that for hydrogen-rich material in the source with a normal period in NGC 6641. The maximum luminosities in the other cluster sources then suggest ultrashort periods in NGC 1851, NGC 6652, NGC 7078-2, and Terzan 2 (Fig. 2).

A third indication comes from the X-ray spectra. A model which is often used – but debated as to its physical correctness – is that in which a sum of blackbody spectra from annuli of the accretion disk is combined with a Comptonized spectrum. The parameters are the inner radius and inner temperature of the disk, and the temperature of the photons entering the Comptonized region, characterized by an optical depth and electron temperature. The best-fit parameters appear to separate the ultrashort systems in NGC 6624 and NGC 6712, for which the inner radius is compatible with a neutron star radius and for which the photons entering the Comptonization region have a temperature compatible with those leaving the disk, from the systems with normal periods in




FIGURE 4. Smoothed grey scale images of NGC 6752 and NGC 6440 as obtained with ROSAT, with the positions of the sources derived from the ROSAT data ($\square$) and the much more numerous positions of the sources detected with Chandra ($\bullet$), illustrating the dramatic increase in resolution and sensitivity. The uncertain direction (‘boresight’) of the ROSAT observation allows a shift between the ROSAT and Chandra image of NGC 6440 by $5''$. ROSAT data from Verbunt & Johnston (2000) and Verbunt et al. (2000); Chandra positions from Pooley et al. (2002a,b).

NGC 6441 and Terzan 6, for which the inner disk radii are too small, and the photons entering the Comptonization region too soft, for the model to be physically viable (Sidoli et al. 2001). This suggests that the sources in NGC 1851, NGC 6652 and Terzan 5 also have an ultrashort period, and those in NGC 6440 and Terzan 2 normal periods (Sidoli et al. 2001, Parmar et al. 2001, Heinke et al. 2003).

Collating these three observational indicators, Verbunt & Lewin (2004) find that the sources in NGC 1851 and NGC 6652 probably, and those in NGC 7078-2 and Terzan 2 possibly, have ultrashort periods (Table 1). The source in NGC 6440 possibly has a normal period. The indications are contradictory in Terzan 2 and absent in NGC 7078-1 and Liller 1. Thus, certainly 2, probably 4, and possibly 6 or 8 of the thirteen bright X-ray sources in globular clusters have ultrashort periods!

VARIOUS TECHNIQUES TO STUDY THE DIM SOURCES

CCDs have a better **spectral resolution** than proportional counters, and as a result the spectral energy distribution of X-ray sources is better determined from observations with Chandra and XMM than with ROSAT (Fig. 1). If a source has a soft spectrum, with a characteristic blackbody temperature of $\lesssim 0.3$ keV say, and an X-ray luminosity $\gtrsim 10^{32} \text{ erg s}^{-1}$, then it is almost certain to be a neutron star accreting from a companion. With this method an increasing number of faint accreting neutron stars has been discovered in globular clusters, showing that the number of such sources is more than ten

times higher than the number of bright sources (Pooley et al. 2003). In general, accreting neutron stars are close to the cluster center, within about 2 core radii, but of order 20% are further from the core. For example, the bright source in NGC 6652 is about six core radii from the center of that cluster (Heinke et al. 2001). With the accuracy of Chandra positions, the error in the position of a source with respect to the cluster center is now dominated by the uncertainty in the cluster center.

The Chandra telescope has a much better **spatial resolution** than ROSAT, and as a result many more faint sources have been discovered in clusters (Figure 4). The improved accuracy of the source positions allows a fairly safe identification of X-ray sources with accurately localized radio pulsars on the basis of positional coincidence alone. A dozen pulsars has thus been detected in 47 Tuc without ambiguity; two more sources are blends of two pulsars each. A single pulsar has been detected in each of the clusters NGC 6397, NGC 6752 and M 28 (Grindlay et al. 2002, D’Amico et al. 2002, Becker et al. 2003). The latter source, already detected with ASCA and ROSAT (Saito et al. 1997, Verbunt 2001), is very bright, and shows that the total X-ray luminosity in the 0.5-2.5 keV range scales directly with the spindown luminosity, as is the case for pulsars (both young and recycled) in the disk of the galaxy (Verbunt et al. 1996).

The **positional accuracy** of Chandra and of the Hubble Space Telescope allows much more secure identification of optical counterparts to the X-ray sources than earlier ROSAT data. To take full advantage of this opportunity, one must align the coordinates from Chandra and HST. From the absolute position of a Chandra source (with accuracy $0.6''$, say) and of the HST objects (with accuracy of $1''$, say; but occasionally as big as $3''$) we must look for counterparts in the HST image in a circle with a radius of $\sqrt{0.6^2 + 1^2} = 1.2''$, or occasionally even larger. On the other hand, once we have securely identified at least one Chandra source with an HST object, we can align the coordinate frames with this match, and the error circles for the remaining sources are much reduced: for sufficiently bright X-ray sources the relative positions are accurate to $\lesssim 0.1''$, the relative optical positions are accurate to $\lesssim 0.1''$, and thus the radius of the search circle now is $\sqrt{0.14^2 + \sigma_m^2}$, where σ_m is the accuracy of the match.

In the case of a rare source type, like pulsars, identifications are secure even with the larger error circles of the absolute astrometry; to be useful for optical identifications, the pulsar radio position must be tied to the optical coordinate frame. A beautiful example of this is the identification of the pulsar in NGC6397 with a Chandra source, on the basis of the binary period, detected both in the radio and in the optical (Ferraro et al. 2001). If such a direct match is not available, one may improve the HST absolute positions by connecting its coordinate system to absolute astrometry. For this purpose, the USNO CCD Astrograph Catalog (UCAC: Zacharias et al. 2000) turns out to be excellent, as shown by Bassa et al. (2004) for NGC 6121. Bassa et al. first identify 91 stars in an ESO 2.2m Wide Field Imager observation with stars in UCAC to determine the coordinates of the ESO WFI image, and then identify about 200 stars in the HST WFPC2 image with stars on the ESO WFI image to determine the coordinates of the HST frame with a 1σ accuracy of $0.12''$ both in right ascension and in declination. This allows identification of three optical objects with Chandra sources, which are then used to align the Chandra frame to the HST frame, with an accuracy of about $0.14''$.

Once we have optical counterparts, we can complete the **classification** of the sources.




FIGURE 5. *Left: Colour – magnitude diagram for NGC6752. Optical counterparts for X-ray sources are indicated $\odot$ and labelled with the Chandra source number; other stars within the X-ray error circles, but presumably not the counterparts are indicated $\square$. From a re-analysis by Bassa (private communication) after Pooley et al. (2002a). Right: The X-ray luminosity as a function of absolute visual magnitude for various source types in 47 Tuc (open symbols) and M4 (filled symbols). The dashed line roughly separates the cataclysmic variables (CV) from the active binaries (AB). Other source types are quiescent low-mass X-ray binaries (qXB), and companions to millisecond pulsars (PSR). Sources which may be either a cataclysmic variable or an active binary are indicated with a star. A 'B' indicates the position of an X-ray source (X2) in M4, which may be a background quasar. The dotted line has L_x hundred times the dashed line. Data from Edmonds et al. (2003), Grindlay et al. (2001) and Bassa et al. (2004).*

From the X-ray data alone we can classify the quiescent low-mass X-ray binaries (from their spectrum and luminosity, see Fig. 1) and the pulsars (from positional coincidence). In the optical colour-magnitude diagram, cataclysmic variables are bluer than the main sequence, and active binaries are more luminous than the main-sequence (Fig. 5). Cataclysmic variables have a higher X-ray to optical flux ratio than magnetic binaries. In a diagram of X-ray luminosity versus absolute visual magnitude, the line

$$\log L_{0.5-2.5\text{keV}}(\text{erg/s}) = -0.4M_V + 33.6$$

roughly separates cataclysmic variables from active binaries (Fig. 5). The variability of cataclysmic variables can cause confusion, if data are combined that were not taken simultaneously. Also, background quasars may masquerade as cataclysmic variables in the L_x - M_V diagram, as may be the case for X2 in M4, tentatively classified as a cataclysmic variable by Bassa et al. (2004), but identical to the blue source, probably a quasar, discovered by Bedin et al. (2003; Bedin & King, private communication). This source is indicated 'B' in Fig. 5. The dotted line, a factor 100 in X-ray luminosity above the dashed line, roughly gives an upper limit to the X-ray luminosities of cataclysmic variables in the galactic disk (based on ROSAT data, Verbunt et al. 1995). Remarkably, two cataclysmic variables in 47 Tuc are well above this line (Grindlay et al. 2001).




FIGURE 6. *Left: The number of X-ray sources detected with Chandra in globular clusters above a threshold limit, as a function of a normalized collision number Γ_n . An arrow indicates a lower limit. The dashed line indicates the best fit: $N \propto \Gamma_n^{0.74}$. After Pooley et al. (2003). Right: central density of globular clusters as a function of core radius, with lines of constant collision number $\Gamma \propto \rho_0^{1.5} r_c^2$ (dotted, steps of factor 10) and of constant destruction number $\gamma \propto \rho_0^{0.5} / r_c$ (dashed, steps of factor 10). Clusters from the left panel are encircled, and (some) indicated with their name. After Verbunt (2003).*

SOME CONSIDERATIONS ON FORMATION RATES

If (quiescent) low-mass X-ray binaries and cataclysmic variables are formed in globular clusters mainly through stellar encounters, their numbers should scale with the collision number Γ of the cluster. Γ scales approximately with the central density ρ_0 and core radius r_c of the cluster as $\Gamma \propto \rho_0^{1.5} r_c^2$ (e.g. Verbunt 2003). Pooley et al. (2003) show that this predictions holds rather well for the X-ray sources with $L_x > 4 \times 10^{30} \text{ erg s}^{-1}$ (in the 0.5-2.5 keV band, Fig. 6). More precisely, with $N \propto \Gamma^\alpha$, they find $\alpha = 0.74 \pm 0.36$, allowing a direct proportionality, but suggesting that N increases slower than Γ . Since most of these sources are actually cataclysmic variables (see e.g. Fig. 5), this indicates that most cataclysmic variables in globular clusters are *not* evolved from primordial binaries, but rather formed via stellar encounters. NGC 6397 is an interesting cluster in this respect, as it has a rather larger number than predicted from its Γ of binaries with neutron stars (one quiescent low-mass X-ray binary, one radio pulsar) and cataclysmic variables. Pooley et al. (2003) suggest that the cluster was more massive in the past, and lost single stars from its outskirts in passages close to the galactic center; binaries escape this stripping process as they reside in the cluster core. The problem with this explanation is that NGC 6397 shows no excess of active binaries in it core. Perhaps the explanation of this is that the active binaries are destroyed in close encounters. The rate at which a single binary undergoes an encounter is roughly $\gamma \propto \rho_0^{0.5} / r_c$ (Verbunt 2003), and NGC 6397 has the highest γ of the globular clusters studied so far (Fig. 6). Since active binaries tend to be wider than cataclysmic variables, they would be more in danger of a destructive encounter.

A wonderful outlook on future work is provided by the study of bright X-ray sources in the globular clusters of M 87: Jordán et al. (2004) show that the probability for a cluster to house a bright X-ray source scales slower with ρ_0 than Γ . They argue that this may be due to destruction of sources in the highest-density clusters.

REFERENCES

1. Bassa, C., Pooley, D., Homer, L., et al. 2004, ApJ 609, 755
2. Becker, W., Swartz, D., Pavlov, G., et al., 2003, ApJ 594, 798
3. Bedin, L., Piotto, G., King, I., and Anderson, J., 2003, AJ 126, 247
4. Chou, Y. and Grindlay, J., 2001, ApJ 563, 934
5. D’Amico, N., Possenti, A., Fici, L., et al. 2002, ApJ 570, L89
6. Deutsch, E., Anderson, S., Margon, B., and Downes, R., 1996, ApJ 472, L97
7. Deutsch, E., Margon, B., and Anderson, S., 2000, ApJ 530, L21
8. Edmonds, P., Gilliland, R., Heinke, C., and Grindlay, J., 2003, ApJ 596, 1177
9. Ferraro, F., Possenti, A., D’Amico, N., and Sabbi, E., 2001, ApJ 561, L93
10. Gendre, B., Barret, D., and Webb, N., 2003, A&A 403, L11
11. Grindlay, J., Heinke, C., Edmonds, P., and Murray, S., 2001, Science 292, 2290
12. Grindlay, J., Camilo, F., Heinke, C., Edmonds, P., Cohn, H., and Lugger, P., 2002, ApJ 581, 470
13. Guainazzi, M., Parmar, A., and Oosterbroek, T., 1999, A&A 349, 819
14. Heinke, C., Edmonds, P., and Grindlay, J., 2001, ApJ 562, 363
15. Heinke, C., Edmonds, P., Grindlay, J., Lloyd, D., Cohn, H., and Lugger, P., 2003, ApJ 590, 809
16. in ’t Zand, J., Hulleman, F., Markwardt, C., et al. 2003, A&A 406, 233
17. Johnston, H., Verbunt, F., and Hasinger, G., 1995, A&A 298, L21
18. Jordán, A., Côté, P., and Ferrarese, L. et al. 2004, ApJ 613, 279
19. King, I., Stanford, S., Albrecht, R., et al., 1993, ApJ 413, L117
20. Kuulkers, E., den Hartog, P., in ’t Zand, J., et al. 2003, A&A 399, 663
21. Parmar, A., Stella, L., and Giommi, P., 1989, A&A 222, 96
22. Parmar, A., Oosterbroek, T., Sidoli, L., Stella, L., and Frontera, F., 2001, A&A 380, 490
23. Podsiadlowski, P., Rappaport, S., and Pfahl, E., 2002, ApJ 565, 1107
24. Pooley, D., Lewin, W., Homer, L., 2002a, ApJ 569, 405
25. Pooley, D., Lewin, W., Verbunt, F., 2002b, ApJ 573, 184
26. Pooley, D., Lewin, W., Anderson, S., et al., 2003, ApJ 591, L131
27. Pylyser, E. and Savonije, G., 1988, A&A 191, 57
28. Rasio, F., Pfahl, E., and Rappaport, S., 2000, ApJ 532, L47
29. Saito, Y., Kawai, N., Kamae, T., et al., 1997, ApJ (Letters) 477, L37
30. Sidoli, L., Parmar, A., Oosterbroek, T., et al. 2001, A&A 368, 451
31. Tutukov, A., Fedorova, A., Ergma, E., and Yungelson, L., 1985, Pis’ma Astron. Zh. 11, 123
32. van der Klis, M., Hasinger, G., Verbunt, F., et al. 1993, A&A 279, L21
33. van der Sluys, M., Verbunt, F., and Pols, O., 2004, A&A in press, astro-ph/0411189
34. van Paradijs, J. and McClintock, J., 1994, A&A 290, 133
35. Verbunt, F., 1987, ApJ 312, L23
36. Verbunt, F., 2001, A&A 368, 137
37. Verbunt, F., 2003, in G. Piotto, G. Meylan, S. Djorgovski, and M. Riello (eds.), New horizons in globular cluster astronomy, pp 245–254, ASP Conf. Ser. 296
38. Verbunt, F. and Johnston, H., 2000, A&A 358, 910
39. Verbunt, F. and Lewin, W., 2004, in W. Lewin and M. van der Klis (eds.), Compact stellar X-ray sources, p. in press, Cambridge University Press
40. Verbunt, F., Bunk, W., Hasinger, G., and Johnston, H., 1995, A&A 300, 732
41. Verbunt, F., Kuiper, L., Belloni, T., Johnston, H., et al. 1996, A&A 311, L9
42. Verbunt, F., van Kerkwijk, M., in ’t Zand, J., and Heise, J., 2000, A&A 359, 960
43. Wijnands, R., Heinke, C., and Grindlay, J., 2002, ApJ 572, 1002
44. Zacharias, N., Urban, S., Zacharias, M., et al., 2000, AJ 120, 2131